



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**MORTALITY—FOREIGN AND INSULAR—COUNTRIES AND CITIES**  
(untabulated).

FRANCE—*Calais*.—Month of March, 1911. Population, 80,000. Total number of deaths from all causes, 111, including measles 5, tuberculosis 24.

GREAT BRITAIN.—Week ended April 1, 1911.

*England and Wales*.—The deaths registered in 77 great towns correspond to an annual rate of 15.4 per 1,000 of the population, which is estimated at 17,160,256.

*Ireland*.—The deaths registered in 21 principal town districts correspond to an annual rate of 20.7 per 1,000 of the population, which is estimated at 1,163,596. The lowest rate was recorded at Wexford, viz, 9.3, and the highest at Tralee, viz, 47.6 per 1,000 of the population.

*Scotland*.—The deaths registered in 8 principal towns correspond to an annual rate of 15.6 per 1,000 of the population, which is estimated at 1,917,875. The lowest was recorded at Aberdeen, viz, 9.4, and the highest at Greenock, viz, 22.5 per 1,000. The total number of deaths from all causes was 573, including diphtheria 3, measles 1, scarlet fever 3.

GREECE—*Patras*.—March 15 to 31, 1911. Population, 39,000. Total number of deaths from all causes, 39, including diphtheria 1, measles 2, tuberculosis 3.

MALTA.—Two weeks ended February 25, 1911. Population, 215,879. Total number of deaths from all causes, 182, including diphtheria 2, tuberculosis 6, typhoid fever 1.

Month of March, 1911. Total number of deaths from all causes reported 233 (deaths for the week ended March 18 not reported), including 20 from tuberculosis.

SIAM—*Bangkok*.—Four weeks ended February 20, 1911. Population, 600,000. Total number of deaths from all causes, not reported. The deaths include smallpox 8, cholera 235, plague 16.

SPAIN—*Almeria*.—Month of March, 1911. Population, 50,910. Total number of deaths from all causes, 139, including diphtheria 1, tuberculosis 14, typhus fever 4.

By authority of the Secretary of the Treasury:

WALTER WYMAN,

Surgeon General,

United States Public Health and Marine-Hospital Service.